

Induced Representations of Quantum Kinematical Algebras and Quantum Mechanics

Oscar Arratia ¹ and Mariano A. del Olmo ²

¹ *Departamento de Matemática Aplicada a la Ingeniería,*

² *Departamento de Física Teórica,*

Universidad de Valladolid, E-47011, Valladolid, Spain

E. mail: oscarr@wmatem.eis.uva.es, olmo@fta.uva.es

February 1, 2008

Abstract

Unitary representations of kinematical symmetry groups of quantum systems are fundamental in quantum theory. We propose in this paper its generalization to quantum kinematical groups. Using the method, proposed by us in a recent paper [1], to induce representations of quantum bicrossproduct algebras we construct the representations of the family of standard quantum inhomogeneous algebras $U_\lambda(iso_\omega(2))$. This family contains the quantum Euclidean, Galilei and Poincaré algebras, all of them in (1+1) dimensions. As byproducts we obtain the actions of these quantum algebras on regular co-spaces that are an algebraic generalization of the homogeneous spaces and q -Casimir equations which play the role of q -Schrödinger equations.

1 Introduction

It is well known the role played by the unitary representations of the (kinematical) symmetry groups of quantum physical systems. One can classify the elementary systems [2] according to them or to obtain their corresponding Schrödinger equations using local representations [3]. In the last years we are involved in a program for constructing a theory of induced representations for quantum algebras [1, 4, 5, 6]. Our aim is to obtain a quantum counterpart of the program that Wigner started in 1939 for Lie groups [2] that has been so fruitful in quantum physics.

Kinematical groups like Poincaré and Galilei, whose physical interest is out of any doubt, have a structure of semidirect product. The quantum version of this kind of structure for quantum groups is that of bicrossproduct [7, 8]. In this case a quantum Lie algebra inherits the ‘semidirect’ structure in the algebra sector and the algebra of functions also has a semidirect product structure in the coalgebra sector. The interest of the quantum versions of the kinematical groups and algebras from a physical point of view is as q -generalizations of the symmetries of the physical space-time in a noncommutative framework. The study of these quantum symmetries and their representations generalizes the Wigner program inside the perspective of the noncommutative geometry [9], whose importance in physics is increasing.

For these groups having structure of semidirect product Mackey’s method [10] provides their unitary representations by induction from the representations of one of their subgroups. In this paper we study the induced representations for some quantum kinematical algebras from a unified point of view. We consider the framework of the Cayley–Klein (CK) pseudo-orthogonal algebras [11] acting in spaces of $(1+1)$ -dimensions, which includes the inhomogeneous algebras like Galilei, Poincaré and Euclidean algebras. The quantum deformation of these CK algebras was performed in [12] (Ref. [13]–[15] present quantum CK algebras in higher dimensions). The above mentioned quantum CK inhomogeneous algebras have a bicrossproduct structure [16] that we profit to obtain their induced representations, given a unified model for all of them.

The induction procedure formulated by us presents a strong algebraic character because we made use of objects like modules, comodules, etc., which, from our point of view, are the appropriate tools to work with the algebraic structures characteristic of the quantum algebras and groups.

In the literature we can find some attempts to develop techniques that generalize the Mackey method [10] of induced representations for semidirect product groups to the quantum case. For instance, Dobrev presented in [17] a method for constructing representations of quantum groups near to ours, since both methods emphasize the dual case, closer to the classical one, and the representations are constructed in the algebra sector. Other authors [18]–[22] have also extended the induction technique to quantum groups but constructing corepresentations, i.e. representations of the coalgebra sector.

The organization of the paper is the following one. In Section 2 we present a mathematical outline of the concepts that we will use along the paper in order to unify notation. Section 3 is devoted to summarize the theory of induced representations of quantum bicrossproduct algebras. We begin to study the induction problem taking into account the deep relation between modules and representations obtaining, in some sense, more deep results from a geometric point of view using the concept of regular co-space. Since our aim is to construct the induced representations of certain quantum inhomogeneous CK algebras Section 4 is devoted to describe briefly these

q -algebras. In Section 5 we construct with some detail the induced representations following the method developed in Section 3. We start with the computation of the flow associated to the action of the generator of one of the factors of the bicrossproduct over the other one. After, we are able to determine the regular co-spaces determining the induced representations, that we obtain below in an explicit way. Moreover, q -Casimir equations are also obtained.

2 Preliminaries

Let $H = (V; m, \eta; \Delta, \epsilon; S)$ be a Hopf algebra with underlying vector space V over the field $\mathbb{K}$ ($\mathbb{C}$ or $\mathbb{R}$), multiplication $m : H \otimes H \rightarrow H$, coproduct $\Delta : H \rightarrow H \otimes H$, unit $\eta : \mathbb{K} \rightarrow H$, counit $\epsilon : H \rightarrow \mathbb{K}$ and antipode $S : H \rightarrow H$.

The algebras involved in this work are infinite dimensional algebras but finitely generated, for this reason we will use a multi-index notation [6]. Let A be an algebra generated by the elements $(a_1, a_2, \dots, a_r)$ such that the ordered monomials $a^n = a_1^{n_1} a_2^{n_2} \dots a_r^{n_r}$ ($n = (n_1, n_2, \dots, n_r) \in \mathbb{N}^r$) form a basis of the linear space underlying to A . An arbitrary product of generators of A is written in a normal ordering if it is expressed in terms of the (ordered) basis $(a^n)_{n \in \mathbb{N}^r}$. The unit of A , 1_A , is denoted by a^0 ($0 \in \mathbb{N}^r$). Multi-factorials and multi-deltas are defined by

$$l! = \prod_{i=1}^n l_i!, \quad \delta_l^m = \prod_{i=1}^n \delta_{l_i}^{m_i}. \quad (2.1)$$

A pairing between two Hopf algebras, H and H' , is a bilinear mapping $\langle \cdot, \cdot \rangle : H \times H' \rightarrow \mathbb{K}$ verifying some defining relations [23]. The pairing is said to be left (right) nondegenerate if $[\langle h, \varphi \rangle = 0, \forall \varphi \in H'] \Rightarrow h = 0$ ($[\langle h, \varphi \rangle = 0, \forall h \in H] \Rightarrow \varphi = 0$). The pairing is said nondegenerate if it is simultaneously left and right nondegenerate. Two Hopf algebras and a nondegenerate pairing, $(H, H', \langle \cdot, \cdot \rangle)$, determine a ‘nondegenerate triplet’. The bases (h^m) of H and (φ_n) of H' are dual with respect to the nondegenerate pairing if

$$\langle h^m, \varphi_n \rangle = c_n \delta_n^m, \quad c_n \in \mathbb{K} - \{0\}. \quad (2.2)$$

Given a nondegenerate triplet and the map $f : H \rightarrow H$. We shall say that the map $f^\dagger : H' \rightarrow H'$, defined by

$$\langle h, f^\dagger(\varphi) \rangle = \langle f(h), \varphi \rangle, \quad (2.3)$$

is the adjoint map to f with respect to $\langle \cdot, \cdot \rangle$.

We will use the endomorphisms of ‘multiplication’ (denoted again by h and φ) and ‘formal derivation’ ($\frac{\partial}{\partial h} \equiv \partial_h$ and $\frac{\partial}{\partial \varphi} \equiv \partial_\varphi$). They verify that $h^\dagger = \partial_\varphi$ and $\varphi^\dagger = \partial_h$. The situation is similar when H and H' are finitely generated. The definition of the formal derivative is

$$\frac{\partial}{\partial h_i} (h_1^{l_1} \dots h_i^{l_i} \dots h_n^{l_n}) = l_i h_1^{l_1} \dots h_i^{l_i-1} \dots h_n^{l_n}. \quad (2.4)$$

The generalization of the ‘multiplication’ operators is not straightforward if the algebras are non commutative. To avoid any confusion the formal operators associated to the generators h_i will be denoted using a bar over the corresponding symbol. The action of these operators is

$$\bar{h}_i (h_1^{l_1} \dots h_i^{l_i} \dots h_n^{l_n}) = h_1^{l_1} \dots h_i^{l_i+1} \dots h_n^{l_n}. \quad (2.5)$$

For the elements $\varphi^i \in H'$ the ‘multiplication’ and derivative operators are defined in a similar way. Note that if H (resp. H') is commutative then $\bar{h}_i$ (resp. $\bar{\varphi}^i$) acts as a multiplication operator, but this is not longer true when the algebra is noncommutative. The adjoint operators are $\bar{h}_i^\dagger = \partial_{\varphi^i}$, $\bar{\varphi}^{i\dagger} = \partial_{h_i}$. The commutation relations for $\bar{h}_i$ and ∂_{h_i} (similar for $\bar{\varphi}^i$ and ∂_{φ^i}) are

$$[\partial_{h_i}, \bar{h}_j] = \delta_{ij}, \quad [\bar{h}_i, \bar{h}_j] = 0, \quad [\partial_{h_i}, \partial_{h_j}] = 0. \quad (2.6)$$

Let (V, α, A) be a triad composed by a unital and associative $\mathbb{K}$ -algebra A , a $\mathbb{K}$ -vector space V and a linear map (called action) $\alpha : A \otimes_{\mathbb{K}} V \rightarrow V$ ($\alpha(a \otimes v) = a \triangleright v$). It is said that (V, α, A) (or $(V, \triangleright, A)$) is a left A -module if:

$$a \triangleright (b \triangleright v) = (ab) \triangleright v, \quad 1 \triangleright v = v, \quad \forall a, b \in A, \quad \forall v \in V. \quad (2.7)$$

Dualizing an A -module a comodule is obtained. The triad $(V, \blacktriangleleft, C)$, where C is an associative $\mathbb{K}$ -coalgebra with counit, V a $\mathbb{K}$ -vector space and $\blacktriangleleft : V \rightarrow C \otimes_{\mathbb{K}} V$ ($v \blacktriangleleft = v^{(1)} \otimes v^{(2)}$) a (linear map) coaction, is said to be a left C -comodule if

$$v^{(1)}_{(1)} \otimes v^{(1)}_{(2)} \otimes v^{(2)} = v^{(1)} \otimes v^{(2)}_{(1)} \otimes v^{(2)}_{(2)}, \quad \epsilon(v^{(1)})v^{(2)} = v, \quad \forall v \in V, \quad (2.8)$$

with $\Delta(c) = c_{(1)} \otimes c_{(2)}$ denoting the coproduct of the elements of C .

A morphism of left A -modules, $(V, \triangleright, A)$ and $(V', \triangleright', A)$, is a linear map, $f : V \rightarrow V'$, equivariant with respect the action, i.e.,

$$f(a \triangleright v) = a \triangleright' f(v), \quad \forall a \in A, \quad \forall v \in V. \quad (2.9)$$

A linear map $f : V \rightarrow V'$ between two C -comodules, $(V, \blacktriangleleft, C)$ and $(V', \blacktriangleleft', C)$ is a morphism if

$$v^{(1)} \otimes f(v^{(2)}) = f(v)^{(1')} \otimes f(v)^{(2')}, \quad \forall v \in V. \quad (2.10)$$

If a bialgebra acts or coacts on a vector space equipped with an additional structure (algebra, coalgebra or bialgebra) some compatibility relations for the action may be demanded [8].

Let A , B , C be an algebra, a bialgebra and a coalgebra, respectively. The left module $(A, \triangleright, B)$ is a B -module algebra if m_A and η_A are morphisms of B -modules. That is, if

$$b \triangleright (aa') = (b_{(1)} \triangleright a)(b_{(2)} \triangleright a'), \quad b \triangleright 1 = \epsilon(b)1, \quad \forall b \in B, \quad \forall a, a' \in A. \quad (2.11)$$

A left B -module $(C, \triangleright, B)$ is a B -module coalgebra if Δ_C and ϵ_C are morphisms of B -modules, i.e., if

$$(b \triangleright c)_{(1)} \otimes (b \triangleright c)_{(2)} = (b_{(1)} \triangleright c_{(1)}) \otimes (b_{(2)} \triangleright c_{(2)}), \quad \epsilon_C(b \triangleright c) = \epsilon_B(b)\epsilon_C(c), \quad \forall b, c \in B.$$

A left B -comodule $(C, \blacktriangleleft, B)$ is said to be a B -comodule coalgebra if Δ_C and ϵ_C are morphisms of B -comodules, i.e.,

$$c^{(1)} \otimes c^{(2)}_{(1)} \otimes c^{(2)}_{(2)} = c_{(1)}^{(1)} c_{(2)}^{(1)} \otimes c_{(1)}^{(2)} \otimes c_{(2)}^{(2)}, \quad c^{(1)} \epsilon_C(c^{(2)}) = (\eta_B \circ \epsilon_C)(c).$$

A left B -comodule $(A, \blacktriangleleft, B)$ is a B -comodule algebra if m_A and η_A are morphisms of B -comodules. Explicitly

$$(aa')_{(1)} \otimes (aa')_{(2)} = a_{(1)} a'_{(1)} \otimes a_{(2)} a'_{(2)}, \quad 1_A \blacktriangleleft = 1_B \otimes 1_A. \quad (2.12)$$

The triad $(B', \triangleright, B)$ is a left B -module bialgebra if simultaneously is a B -module algebra and a B -module coalgebra; $(B', \blacktriangleleft, B)$ is a left B -comodule bialgebra if is a B -comodule algebra and a B -comodule coalgebra.

A regular module (comodule) is an A -module (C -comodule) whose vector space is the underlying vector space of the algebra A (coalgebra C). The action (coaction) is defined in terms of the algebra product (coalgebra coproduct). If B is a bialgebra, the regular B -module $(B, \triangleright, B)$, whose regular action is $b \triangleright b' = bb'$, is a module coalgebra. The regular module $(B^*, \triangleleft, B)$, obtained by dualization, is a module algebra with regular action $\varphi \triangleleft b = \langle \varphi_{(1)}, b \rangle \varphi_{(2)}$ with $b \in B$, $\varphi \in B^*$.

Let K and L be two Hopf algebras, with $(L, \triangleleft, K)$ a right K -module algebra and $(K, \blacktriangleleft, L)$ a left L -comodule coalgebra. The tensor product $K \otimes L$ is equipped simultaneously with the semidirect structures of algebra $K \bowtie L$ and coalgebra $K \blacktriangleright L$. If some compatibility conditions are verified $K \bowtie L$ and $K \blacktriangleright L$ determine a Hopf algebra called (right-left) bicrossproduct and denoted by $K \blacktriangleright L$ [7, 8].

Let $\mathcal{K}$ and $\mathcal{L}$ be two Hopf algebras and $(\mathcal{L}, \triangleright, \mathcal{K})$ and $(\mathcal{K}, \blacktriangleright, \mathcal{L})$ a left K -module algebra and a right L -comodule coalgebra, respectively, verifying the corresponding compatibility conditions. Then $\mathcal{L} \bowtie \mathcal{K}$ and $\mathcal{L} \blacktriangleright \mathcal{K}$ determine a Hopf algebra, $\mathcal{L} \blacktriangleright \mathcal{K}$, called (left-right) bicrossproduct.

These two bicrossproduct structures are related by duality: if K and L are finite dimensional bialgebras and the K -module algebra $(L, \triangleleft, K)$ and the L -comodule coalgebra $(K, \blacktriangleleft, L)$ verify the compatibility conditions determining the bicrossproduct $K \blacktriangleright L$, then $(K \blacktriangleright L)^* = K^* \blacktriangleright L^*$.

It has been proved in Ref. [1] that dual bases and $*$ -structures over bicrossproduct Hopf algebras may be constructed when the corresponding bases and $*$ -structures of the bicrossproduct the factors are known. Thus, let $H = K \blacktriangleright L$ be a bicrossproduct Hopf algebra and $(K, K^*, \langle \cdot, \cdot \rangle_1)$ and $(L, L^*, \langle \cdot, \cdot \rangle_2)$ nondegenerate triplets, then the expression

$$\langle kl, \kappa \lambda \rangle = \langle k, \kappa \rangle_1 \langle l, \lambda \rangle_2. \quad (2.13)$$

defines a nondegenerate pairing between H and H^* . If (k_m) and (κ_m) are dual bases for K and K^* , and (l_n) and (λ_n) for L and L^* , then $(k_m l_n)$ and $(\kappa^m \lambda^n)$ are dual bases for H and H^* .

It can be proved that given a bicrossproduct Hopf algebra, $H = K \blacktriangleright L$, such that K and L are equipped with $*$ -structures verifying the compatibility relation $(l \triangleleft k)^* = l^* \triangleleft S(k)^*$, there is a $*$ -structure on the algebra sector of H is determined by

$$(kl)^* = l^* k^*, \quad k \in K, l \in L. \quad (2.14)$$

3 Induced representations of bicrossproduct algebras

The main results about the theory of induced representations for quantum bicrossproduct algebras will be summarized in this Section (see Refs. [1, 4, 5, 6] for a complete description of induced representations of quantum algebras).

As we have seen in the previous Section different actions are involved, henceforth we will denote them by the following symbols (or their symmetric for the corresponding right actions and coactions): $\triangleright$ ($\blacktriangleleft$) (bicrossproduct actions (coactions)), $\vdash$ (induced and inducing representations) and $\succ$, $\prec$ (regular actions).

Let $(H, \mathcal{H}, \langle \cdot, \cdot \rangle)$ be a nondegenerate triplet and L a commutative subalgebra of H . Suppose that $\{l_1, \dots, l_s\}$ is a system of generators of L which is completed with $\{k_1, \dots, k_r\}$ to get a system of generators of H , such that $(l_n)_{n \in \mathbb{N}^s}$ is a basis of L and $(k_m l_n)_{(m,n) \in \mathbb{N}^r \times \mathbb{N}^s}$ a basis of H . Moreover, there is a generator system of $\mathcal{H}$, $\{\kappa_1, \dots, \kappa_r, \lambda_1, \dots, \lambda_s\}$, such that $(\kappa^m \lambda^n)_{(m,n) \in \mathbb{N}^r \times \mathbb{N}^s}$ is a basis of $\mathcal{H}$ dual of that of H with pairing

$$\langle k_m l_n, \kappa^{m'} \lambda^{n'} \rangle = m! n! \delta_m^{m'} \delta_n^{n'}. \quad (3.1)$$

In order to construct the representation of H induced by the character of L , determined by $a = (a_1, \dots, a_s) \in \mathbb{K}^s$ (explicitly, $1 \dashv l_n = a_n = a_1^{n_1} \dots a_s^{n_s}$, $n \in \mathbb{N}^s$) we need to know its carrier space $\mathbb{K}^\uparrow$ and the action of H on it. The elements of $\mathbb{K}^\uparrow$ are those of $\text{Hom}_{\mathbb{K}}(H, \mathbb{K})$ verifying the invariance condition

$$f(hl) = f(h) \dashv l, \quad \forall l \in L, \quad \forall h \in H. \quad (3.2)$$

They can be written as

$$f = \sum_{(m,n) \in \mathbb{N}^r \times \mathbb{N}^s} f_{mn} \kappa^m \lambda^n \quad (3.3)$$

by identifying $\mathbb{K}^\uparrow = \text{Hom}_{\mathbb{K}}(H, \mathbb{K})$ with $\mathcal{H}$ via the pairing. The equivariance condition (3.2)

$$\langle hl, f \rangle = \langle h, f \rangle \dashv l, \quad \forall l \in L, \quad \forall h \in H, \quad (3.4)$$

together with duality give the relation

$$m! n! f_{mn} = \langle k_m l_n, f \rangle = \langle k_m, f \rangle a_n = m! f_{m0} a_n. \quad (3.5)$$

Hence, the elements of $\mathbb{K}^\uparrow$ are

$$f = \kappa \psi, \quad \kappa \in \mathcal{K}, \quad \psi = e^{a_1 \lambda_1} \dots e^{a_s \lambda_s}, \quad (3.6)$$

where $\mathcal{K}$ is the subspace of $\mathcal{H}$ generated by the linear combinations of the ordered monomials $(\kappa^m)_{m \in \mathbb{N}^r}$. Because ψ is product of exponentials, $\mathcal{K}$ and $\mathbb{K}^\uparrow$ are isomorphic ($\kappa \rightarrow \kappa \psi$).

The action of H on $\mathbb{K}^\uparrow$ is determined knowing the action over the basis elements $(\kappa^p \psi)_{p \in \mathbb{N}^r}$. Putting

$$(\kappa^p \psi) \dashv h = \sum_{(m,n) \in \mathbb{N}^r \times \mathbb{N}^s} [h]_{mn}^p \kappa^m \lambda^n, \quad p \in \mathbb{N}^r, \quad (3.7)$$

the coefficients $[h]_{mn}^p$ are evaluated by means of duality

$$m! n! [h]_{mn}^p = \langle (\kappa^p \psi) \dashv h, k_m l_n \rangle = \langle \kappa^p \psi, h k_m l_n \rangle = \langle \kappa^p \psi, h k_m \rangle a_n. \quad (3.8)$$

The properties of the action allow to compute it only for the generators of H instead of considering an arbitrary element of H . The problem reduces to write $h k_m$ in normal ordering to get the value of the paring in (3.8). Since in many cases this task is very cumbersome, our objective will be to take advantage of the bicrossproduct structure to simplify the computations.

In the following we will restrict ourselves to Hopf algebras having a bicrossproduct structure like $H = \mathcal{K} \bowtie \mathcal{L}$, such that $\mathcal{K}$ is cocommutative and $\mathcal{L}$ commutative. Let us suppose that the algebras $\mathcal{K}$ and $\mathcal{L}$ are finite generated by the sets $\{k_i\}_{i=1}^r$ and $\{l_i\}_{i=1}^s$, respectively, the k_i 's are primitive and $(k_n)_{n \in \mathbb{N}^r}$ and $(l_m)_{m \in \mathbb{N}^s}$ are bases of the vector spaces underlying to $\mathcal{K}$ and $\mathcal{L}$, respectively. Let $\mathcal{K}^*$ and $\mathcal{L}^*$ be the dual algebras of $\mathcal{K}$ and $\mathcal{L}$ having dual systems to those of $\mathcal{K}$ and $\mathcal{L}$ with analogue properties to them. So, duality between H and H^* is given by (3.1).

We are interested in the construction of the representations induced by 'real' characters of the commutative sector $\mathcal{L}$. We will show that the solution of this problem can be reduced to the study of certain dynamical systems which present, in general, a nonlinear action. The above results can be summarized in the following theorem.

Theorem 3.1. *The carrier space of the representation of H induced by the character a of $\mathcal{L}$*

$$1 \dashv l_n = a_n, \quad a \in \mathbb{C}^s, \quad n \in \mathbb{N}^s, \quad (3.9)$$

is isomorphic to $\mathcal{K}^*$ and is constituted by the elements of the form

$$\kappa \psi, \quad \kappa \in \mathcal{K}^*, \quad \psi = e^{a_1 \lambda_1} e^{a_2 \lambda_2} \dots e^{a_s \lambda_s}, \quad \lambda_i \in \mathcal{L}^*. \quad (3.10)$$

The induced action of the elements of H over the elements of the carrier space $\mathbb{C}^\uparrow$ is given by

$$f \dashv h = \sum_{m \in \mathbb{N}^r} \kappa^m \langle h \frac{k_m}{m!}, f \rangle \psi, \quad f \in \mathbb{C}^\uparrow. \quad (3.11)$$

To obtain the explicit action of the generators of $\mathcal{K}$ and $\mathcal{L}$ in the induced representation, let us start identifying $\mathcal{L}$, because is commutative, with the algebra of functions $F(\mathbb{R}^s)$ by the morphism $\mathcal{L} \rightarrow F(\mathbb{R}^s)$, $(l \mapsto \tilde{l})$, mapping the generators of $\mathcal{L}$ into the canonical projections

$$\tilde{l}_j(x) = x_j, \quad \forall x = (x_1, x_2, \dots, x_s) \in \mathbb{R}^s, \quad j = 1, 2, \dots, s. \quad (3.12)$$

This identification allows to chose a $*$ -structure keeping invariant the generators of $\mathcal{L}$ by

$$l_j^* = l_j, \quad j = 1, 2, \dots, s. \quad (3.13)$$

Hence, the characters of $\mathcal{L}$ (3.9) compatible with (3.13) are real. They can be written now as

$$1 \dashv l = \tilde{l}(a), \quad a \in \mathbb{R}^s. \quad (3.14)$$

Also, the right action of $\mathcal{K}$ on $\mathcal{L}$ can be carried to $F(\mathbb{R}^s)$. Since the generators of $\mathcal{K}$ are primitive, they act by derivations on the $\mathcal{K}$ -module algebra of $\mathcal{K} \bowtie \mathcal{L}$ inducing vector fields, X_i , on $\mathbb{R}^s$ by

$$X_i \tilde{l} = \widetilde{l \triangleleft k_i}, \quad i = 1, 2, \dots, r. \quad (3.15)$$

The associated flow to X_i , $\Phi_i : \mathbb{R} \times \mathbb{R}^s \rightarrow \mathbb{R}^s$, is given by

$$(X_i f)(x) = (Df_{x, \Phi_i})(0), \quad (3.16)$$

where $f_{x, \Phi_i}(t) = f \circ \Phi_i^t(x)$ and D is the derivative operator. Thus, we can state the following theorem that allows us to have explicit formulae for the action.

Theorem 3.2. *The explicit action of the generators of $\mathcal{K}$ and $\mathcal{L}$ in the induced representation determined in Theorem 3.1 and realized in the space $\mathcal{K}^*$ is given by the following expressions:*

$$\begin{aligned}\kappa \dashv k_i &= \kappa \prec k_i, \\ \kappa \dashv l_j &= \kappa \hat{l}_j \circ \Phi_{(\kappa_1, \kappa_2, \dots, \kappa_r)}(a),\end{aligned}\tag{3.17}$$

where $i \in \{1, \dots, r\}$, $j \in \{1, \dots, s\}$, the symbol $\prec$ denotes the regular action of $\mathcal{K}$ on $\mathcal{K}^*$ and $\Phi_{(\kappa_1, \kappa_2, \dots, \kappa_r)} = \Phi_r^{\kappa_r} \circ \dots \circ \Phi_2^{\kappa_2} \circ \Phi_1^{\kappa_1}$.

We can reformulate the induction procedure in terms of modules because their deep relation with representations [1, 6]. The regular H -modules associated to a Hopf algebra H and its dual H^* , $(H, \prec, H)$, $(H^*, \succ, H)$, $(H, \succ, H)$ and $(H^*, \prec, H)$, will help us to describe the actions as well as the carrier spaces involved in the induced representations. The modules $(H^*, \succ, H)$ and $(H^*, \prec, H)$ are called co-spaces since they can be considered as an algebraic generalization of the concept of G -space.

The commutative or cocommutative Hopf algebras are, in fact, of the form $F(G)$ or $\mathbb{C}[G]$, the group algebra of G , (or $U(\mathfrak{g})$) for any group G [8]. Hence, the bicrossproduct Hopf algebras, that we consider in this paper, can be factorized as

$$H = U(\mathfrak{k}) \bowtie F(L), \quad H^* = F(K) \bowtie U(\mathfrak{l}).\tag{3.18}$$

where K and L are Lie groups with associated Lie algebras $\mathfrak{k}$ and $\mathfrak{l}$, respectively.

The use of elements of H and H^* , like

$$\begin{aligned}k\lambda &\in H, & k &\in K, & \lambda &\in F(L), \\ \kappa l &\in H^*, & \kappa &\in F(K), & l &\in L,\end{aligned}\tag{3.19}$$

instead of the standard bases of ordered monomials, allows an effective description of the H -regular modules.

Theorem 3.3. *The action on each of the four regular H -modules is:*

$$\begin{aligned}(H, \prec, H) &: (k\lambda) \prec k' = kk'(\lambda \triangleleft k'), & (k\lambda) \prec \lambda' &= k\lambda\lambda'; \\ (H^*, \succ, H) &: k' \succ (\kappa l) = (k' \succ \kappa)(k' \triangleright l), & \lambda' \succ (\kappa l) &= \lambda'(l)\kappa l; \\ (H, \succ, H) &: k' \succ (k\lambda) = k'k\lambda, & \lambda' \succ (k\lambda) &= k(\lambda' \triangleleft k)\lambda; \\ (H^*, \prec, H) &: (\kappa l) \prec k' = (\kappa \prec k')l, & (\kappa l) \prec \lambda' &= \kappa(\lambda' \circ \hat{l})l;\end{aligned}\tag{3.20}$$

where $\hat{l}$ is the map, $K \rightarrow L$ ($k \mapsto k \triangleright l$), projecting the group K on the orbit passing through $l \in L$ such that $\langle l^{(1)}, \lambda \rangle l^{(2)} = \lambda \circ \hat{l}$, for all $k, k' \in K$, $\lambda, \lambda' \in F(L)$, $\kappa \in F(K)$ and $l \in L$.

Notice that in this theorem we have taken into account that in $(U(\mathfrak{l}), \succ, F(L))$ holds

$$\lambda \succ l = \lambda(l)l, \quad \forall \lambda \in F(L), \quad \forall l \in L.\tag{3.21}$$

Now we can carry a complete analysis of the representations of $H = U(\mathfrak{k}) \bowtie F(L)$ induced by the one-dimensional modules of its commutative sector. Note that the set of characters of

the algebra $F(L)$ is its spectrum and, hence, the spectrum of $F(L)$ is isomorphic to L . Fixed an element l of L , the character is given by

$$1 \dashv \lambda = \lambda(l), \quad \lambda \in F(L). \quad (3.22)$$

The carrier space $\mathbb{C}^\uparrow$ of the representation of $H = U(\mathfrak{k}) \bowtie F(L)$ induced by (3.22) is the set of elements $f \in H^*$ satisfying the condition

$$\lambda \succ f = \lambda(l)f, \quad \forall \lambda \in F(L). \quad (3.23)$$

Expanding f in terms of the bases of $\mathfrak{k}$ and $\mathfrak{l}$, imposing the equivariance condition and considering the definition of second kind coordinates λ_j over the group L we obtain that

$$f = \kappa l, \quad \kappa \in F(K). \quad (3.24)$$

The right regular action determines the action on the induced module that can be carried to $F(K)$ using the isomorphism $F(K) \rightarrow \mathbb{C}^\uparrow$ ($\kappa \mapsto \kappa l$). So,

$$\kappa \dashv k = \kappa \prec k, \quad \kappa \dashv \lambda = \kappa(\lambda \circ \hat{l}). \quad (3.25)$$

Comparing these expressions with those of Theorem 3.2 we see that the action of $U(\mathfrak{k})$ is determined by the regular action. The action of $F(L)$ is multiplicative and its evaluation is essentially reduced to obtain the flows associated to the action of K on L derived from the bicrossproduct structure of H .

Finally, next theorem summarizes the induction procedure for bicrossproduct algebras.

Theorem 3.4. *Let us consider an element $l \in L$ supporting a global action of the group K . The carrier space, $\mathbb{C}^\uparrow$, of the representation of H induced by the character determined by l is the set of elements of H^* of the form*

$$\kappa l, \quad \kappa \in F(K). \quad (3.26)$$

There is an isomorphism between $\mathbb{C}^\uparrow$ and $F(K)$ given by the map $\kappa \mapsto \kappa l$. The action induced by the elements of the form $k \in K$ and $\lambda \in F(L)$ in the space $F(K)$ is

$$\begin{aligned} \kappa \dashv k &= \kappa \prec k \\ \kappa \dashv \lambda &= \kappa(\lambda \circ \hat{l}). \end{aligned} \quad (3.27)$$

The modules induced by l and $k \triangleright l$ are isomorphic. So, the induction algorithm establishes a correspondence between the space of orbits L/K and the set of equivalence classes of representations.

3.1 Local representations

The quantum counterpart of the local representations [3] of Lie groups can be obtained inducing from representations of the subalgebra $U(\mathfrak{k})$. Given a character κ of $U(\mathfrak{k})$,

$$\kappa \in \text{Spectrum } U(\mathfrak{k}) \subset F(K), \quad k \vdash 1 = \kappa(k), \quad (3.28)$$

the carrier space, $\mathbb{C}^\dagger$, of the representation induced by κ is determined by the equivariance condition

$$f \prec k = \kappa(k)f, \quad \forall k \in U(\mathfrak{k}) \quad (3.29)$$

obtaining that the elements of $\mathbb{C}^\dagger$ are of the form

$$f = \kappa l, \quad l \in U(\mathfrak{l}). \quad (3.30)$$

The isomorphism $U(\mathfrak{l}) \rightarrow \mathbb{C}^\dagger$ ($l \mapsto \kappa l$), allows to realize the induced representation over $U(\mathfrak{l})$

$$k \vdash l = \kappa(k) k \triangleright l, \quad \lambda \vdash l = \lambda(l)l. \quad (3.31)$$

4 Quantum $\mathfrak{iso}_\omega(2)$ algebras

The called CK pseudo-orthogonal algebras $\mathfrak{so}_{\omega_1, \omega_2, \dots, \omega_N}(N+1)$ is a family of $(N+1)N/2$ dimensional real Lie algebras characterized by N real parameters $(\omega_1, \omega_2, \dots, \omega_N)$ [11, 16]. In the ‘geometric’ basis $(J_{ij})_{0 \leq i < j \leq N}$ the nonvanishing commutators are

$$[J_{ij}, J_{ik}] = \omega_{ij} J_{jk} \quad [J_{ij}, J_{jk}] = -J_{ik} \quad [J_{ik}, J_{jk}] = \omega_{jk} J_{ij}, \quad (4.1)$$

with $0 < i < j < k < N$ and $\omega_{ij} = \prod_{s=i+1}^j \omega_s$.

The parameters ω_i , in fact, only take the values 1, 0 and -1 since the generators J_{ij} can be rescaled. When $\omega_i \neq 0 \forall i$, the Lie algebra $\mathfrak{so}_{\omega_1, \omega_2, \dots, \omega_N}(N+1)$ is isomorphic to some of the pseudo-orthogonal algebras $\mathfrak{so}(p, q)$ with $p + q = N + 1$ and $p \geq q \geq 0$. If some of the ω_i vanish the corresponding algebra is inhomogeneous.

In this paper we are interested in the particular case of $N = 2$ and $\omega_1 = 0$. These inhomogeneous algebras $\mathfrak{so}_{0, \omega_2}(3)$ can be realized as algebras of groups of affine transformations on $\mathbb{R}^2$ [11] (i.e, the Euclidean group when $\omega_2 = 1$, the Galilei group when $\omega_2 = 0$ and the Poincaré group when $\omega_2 = -1$). In this case the generators J_{0i} are denoted by P_i stressing their role as generators of translations. The remaining generator $J_{12} \equiv J$ is associated to compact ($\omega_2 = 1$) and noncompact rotations (galilean and lorentzian boosts).

The algebra $\mathfrak{so}_{0, \omega_2}(3)$, that we shall denote by $\mathfrak{iso}_\omega(2)$, is characterized by the commutators

$$[J, P_1] = P_2, \quad [J, P_2] = -\omega P_1, \quad [P_1, P_2] = 0. \quad (4.2)$$

We shall use the following ‘generalized trigonometric functions’ [11, 12]

$$C_\omega(x) = \frac{e^{\sqrt{-\omega}x} + e^{-\sqrt{-\omega}x}}{2}, \quad S_\omega(x) = \frac{e^{\sqrt{-\omega}x} - e^{-\sqrt{-\omega}x}}{2\sqrt{-\omega}}. \quad (4.3)$$

When $\omega < 0$ ($\omega > 0$) these expressions become the trigonometric (hyperbolic) functions. For $\omega = 0$ the parabolic functions $C_0(x) = 1$ and $S_0(x) = x$ are obtained. The functions (4.3) satisfy

identities similar to those of the usual trigonometric functions. Some useful properties that will be used in next computations are:

$$\begin{aligned} C_\omega^2(x) + \omega S_\omega^2(x) &= 1, \\ C_\omega(x+y) &= C_\omega(x)C_\omega(y) - \omega S_\omega(x)S_\omega(y), \quad S_\omega(x+y) = S_\omega(x)C_\omega(y) + C_\omega(x)S_\omega(y), \\ C'_\omega(x) &= -\omega S_\omega(x), \quad S'_\omega(x) = C_\omega(x). \end{aligned} \quad (4.4)$$

A simultaneous standard deformation for all the enveloping algebras $U(\mathfrak{so}_{\omega_1, \omega_2}(3))$ was introduced in [12], The particular case of the deformed Hopf algebras $U_\lambda(\mathfrak{iso}_\omega(2))$ is obtained taking $\omega_1 = 0$ and, obviously, $\omega_2 = \omega$.

It was proved in [16] that the standard quantum Hopf algebras $U_\lambda(\mathfrak{iso}_{\omega_2, \omega_3, \dots, \omega_N}(N))$ have a structure of bicrossproduct. So, $U_\lambda(\mathfrak{iso}_\omega(2))$ is characterized in a basis adapted to its bi-crossproduct structure by

$$\begin{aligned} [J, P_1] &= \frac{1-e^{-2\lambda P_2}}{2\lambda} + \frac{1}{2}\lambda\omega P_1^2, \quad [J, P_2] = -\omega P_1; \\ \Delta(P_1) &= P_1 \otimes 1 + e^{-\lambda P_2} \otimes P_1, \quad \Delta(P_2) = P_2 \otimes 1 + 1 \otimes P_2, \quad \Delta(J) = J \otimes 1 + e^{-\lambda P_2} \otimes J; \\ \epsilon(P_1) &= \epsilon(P_2) = \epsilon(J) = 0; \\ S(P_1) &= -e^{\lambda P_2} P_1, \quad S(P_2) = -P_2, \quad S(J) = -e^{\lambda P_2} J. \end{aligned}$$

The bicrossproduct structure $U_\lambda(\mathfrak{iso}_\omega(2)) = \mathcal{K} \bowtie \mathcal{L}$, where $\mathcal{L}$ is the Hopf subalgebra spanned by (P_1, P_2) and $\mathcal{K}$ is the commutative and cocommutative Hopf algebra generated by J , is determined by the action of $\mathcal{K}$ on $\mathcal{L}$ given by

$$P_1 \triangleleft J = [P_1, J] = -\left[\frac{1-e^{-2\lambda P_2}}{2\lambda} + \frac{1}{2}\lambda\omega P_1^2\right], \quad P_2 \triangleleft J = [P_2, J] = \omega P_1, \quad (4.5)$$

and the left coaction of $\mathcal{L}$ on $\mathcal{K}$, which over the generator of $\mathcal{K}$ takes the value

$$J \blacktriangleleft = e^{-\lambda P_2} \otimes J. \quad (4.6)$$

The dual algebra $F_\lambda(ISO_\omega(2))$ is generated by the local coordinates φ, a_1, a_2 . Its commutators, coproduct, counit and antipode are given by

$$\begin{aligned} [a_1, \varphi] &= \lambda(1 - C_\omega(\varphi)), \quad [a_2, \varphi] = \lambda S_\omega(\varphi), \quad [a_1, a_2] = \lambda a_1; \\ \Delta(a_1) &= a_1 \otimes C_\omega(\varphi) + a_2 \otimes \omega S_\omega(\varphi) + 1 \otimes a_1, \\ \Delta(a_2) &= -a_1 \otimes S_\omega(\varphi) + a_2 \otimes C_\omega(\varphi) + 1 \otimes a_2, \\ \Delta(\varphi) &= \varphi \otimes 1 + 1 \otimes \varphi; \\ \epsilon(a_1) &= \epsilon(a_2) = \epsilon(\varphi) = 0; \end{aligned} \quad (4.7)$$

$$S(a_1) = -C_\omega(\varphi)a_1 - \omega S_\omega(\varphi)a_2, \quad S(a_2) = S_\omega(\varphi)a_1 - C_\omega(\varphi)a_2, \quad S(\varphi) = -\varphi.$$

This Hopf algebra exhibits the bicrossproduct structure $F_\lambda(ISO_\omega(2)) = \mathcal{K}^* \blacktriangleright \mathcal{L}^*$, dual of the above one, with $\mathcal{K}^*$ generated by φ and $\mathcal{L}^*$ by a_1, a_2 . The left action of $\mathcal{L}^*$ over $\mathcal{K}^*$ is given by

$$a_1 \triangleright \varphi = \lambda(1 - C_\omega(\varphi)), \quad a_2 \triangleright \varphi = \lambda S_\omega(\varphi). \quad (4.8)$$

The right coaction of $\mathcal{K}^*$ over $\mathcal{L}^*$ takes the following values over the generators of $\mathcal{L}^*$

$$\blacktriangleright a_1 = a_1 \otimes C_\omega(\varphi) + a_2 \otimes \omega S_\omega(\varphi), \quad \blacktriangleright a_2 = -a_1 \otimes S_\omega(\varphi) + a_2 \otimes C_\omega(\varphi). \quad (4.9)$$

The results mentioned in the last paragraph of Section 2 about dual bases and $*$ -structures over bicrossproduct Hopf algebras allow to construct a pair of dual bases in such a way that the duality form between $U_\lambda(\mathfrak{iso}_\omega(2))$ and $F_\lambda(ISO_\omega(2))$ is

$$\langle J^m P_1^n P_2^p, \varphi^q a_1^r a_2^s \rangle = m!n!p! \delta_m^q \delta_n^r \delta_p^s. \quad (4.10)$$

5 Representations of $U_\lambda(\mathfrak{iso}_\omega(2))$

Firstly, according to the theory of representations of bicrossproduct algebras displayed in Section 3 and, in particular, Theorem 3.2, we need to know the flow associated to the action of $\mathcal{K}$ on $\mathcal{L}$.

5.1 One-parameter flow for $U_\lambda(\mathfrak{iso}_\omega(2))$

In Ref. [16] the factor $\mathcal{K}$ is interpreted as the enveloping algebra $U(\mathfrak{so}_\omega(2))$, while $\mathcal{L}$ is seen as a deformation of the algebra of the group of the translations in the plane T_2 , $U_\lambda(\mathfrak{t}_2)$. Hence,

$$U_\lambda(\mathfrak{iso}_\omega(2)) = U(\mathfrak{so}_\omega(2)) \blacktriangleright U_\lambda(\mathfrak{t}_2), \quad F_\lambda(ISO_\omega(2)) = F(SO_\omega(2)) \blacktriangleright F_\lambda(T_2). \quad (5.1)$$

However, the interpretation given here by us is different since

$$U_\lambda(\mathfrak{iso}_\omega(2)) = U(\mathfrak{so}_\omega(2)) \blacktriangleright F(T_{\lambda,2}), \quad (5.2)$$

being $T_{\lambda,2}$ the Lie group with composition law

$$(\alpha'_1, \alpha'_2)(\alpha_1, \alpha_2) = (\alpha'_1 + e^{-\lambda\alpha'_2}\alpha_1, \alpha'_2 + \alpha_2). \quad (5.3)$$

Now the generators P_1 and P_2 are considered as a global coordinate system over $T_{\lambda,2}$, i.e.,

$$P_1(\alpha_1, \alpha_2) = \alpha_1, \quad P_2(\alpha_1, \alpha_2) = \alpha_2. \quad (5.4)$$

The module algebra structure included in $U_\lambda(\mathfrak{iso}_\omega(2)) = U(\mathfrak{so}_\omega(2)) \blacktriangleright F(T_{\lambda,2})$ determines the action of $SO_\omega(2)$ over $T_{\lambda,2}$ by means of the vector field

$$\hat{J}_{\omega,\lambda} = - \left[\frac{1 - e^{-2\lambda P_2}}{2\lambda} + \frac{1}{2} \lambda \omega P_1^2 \right] \frac{\partial}{\partial P_1} + \omega P_1 \frac{\partial}{\partial P_2}. \quad (5.5)$$

Prior to determine the flow associated to $\hat{J}_{\omega,\lambda}$ we can obtain a first qualitative information by identifying the fixed points of $\hat{J}_{\omega,\lambda}$. When $\omega \neq 0$ only the origin $(0,0)$ is an equilibrium point,

but if $\omega = 0$ the set of fixed points is the straight line of equation $\alpha_2 = 0$. The deformation due to λ does not change the character of these points. Thus, the point is elliptic if $\omega > 0$ and hyperbolic if $\omega < 0$.

The one-form

$$\eta_\rho = \rho \left[\omega P_1 dP_1 + \left(\frac{1 - e^{-2\lambda P_2}}{2\lambda} + \frac{1}{2} \lambda \omega P_1^2 \right) dP_2 \right], \quad \rho \in F(T_{\lambda,2}), \quad (5.6)$$

verifies $\hat{J}_{\omega,\lambda} \rfloor \eta_\rho = 0$. Taking $\rho_0 = \lambda^2 e^{\lambda P_2}$ one obtains that η_{ρ_0} is exact and invariant under $\hat{J}_{\omega,\lambda}$. Adding a constant and rescaling the invariant function under the action of J

$$h = \frac{1}{2} \omega \lambda^2 P_1^2 e^{\lambda P_2} + \cosh(\lambda P_2), \quad (5.7)$$

we obtain a central element of $U_\lambda(\mathfrak{iso}_\omega(2))$

$$C_{\omega,\lambda} = \omega P_1^2 e^{\lambda P_2} + 2 \frac{\cosh(\lambda P_2) - 1}{\lambda^2} = \omega P_1^2 e^{\lambda P_2} + \frac{4}{\lambda^2} \sinh^2\left(\frac{\lambda}{2} P_2\right) \quad (5.8)$$

such that in the limit $\lambda \rightarrow 0$ we recover the nondeformed Casimir $C_{\omega,0} = \omega P_1^2 + P_2^2$ of $\mathfrak{iso}_\omega(2)$.

The computation of the trajectories of $\hat{J}_{\omega,\lambda}$ requires to solve the equation system

$$\begin{aligned} \dot{\alpha}_1 &= - \left[\frac{1 - e^{-2\lambda \alpha_2}}{2\lambda} + \frac{1}{2} \lambda \omega \alpha_1^2 \right], \\ \dot{\alpha}_2 &= \omega \alpha_1. \end{aligned} \quad (5.9)$$

The integral curve γ can be expressed as

$$\begin{aligned} \alpha_1(t) &= - \frac{1}{\lambda} \frac{\sinh(\lambda \beta) S_\omega(t)}{\cosh(\lambda \beta) + \sinh(\lambda \beta) C_\omega(t)}, \\ \alpha_2(t) &= \frac{1}{\lambda} \ln[\cosh(\lambda \beta) + \sinh(\lambda \beta) C_\omega(t)]. \end{aligned} \quad (5.10)$$

Using the fact that $\Phi_{\omega,\lambda}^t(\gamma(\tau)) = \gamma(\tau + t)$ the flow can be evaluated

$$\begin{aligned} \Phi_{\omega,\lambda}^t(\alpha_1, \alpha_2) &= \\ &\left(\frac{2\alpha_1 \lambda e^{\lambda \alpha_2} C_\omega(t) + (\omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2} - \sinh(\lambda \alpha_2)) S_\omega(t)}{2\lambda \cosh(\lambda \alpha_2) + \omega \lambda^3 \alpha_1^2 e^{\lambda \alpha_2} + (2\lambda \sinh(\lambda \alpha_2) - \omega \lambda^3 \alpha_1^2 e^{\lambda \alpha_2}) C_\omega(t) + 2\omega \lambda^2 \alpha_1 e^{\lambda \alpha_2} S_\omega(t)}, \right. \\ &\left. \frac{1}{\lambda} \ln \left[\cosh(\lambda \alpha_2) + \frac{1}{2} \omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2} + (\sinh(\lambda \alpha_2) - \frac{1}{2} \omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2}) C_\omega(t) + \omega \lambda \alpha_1 e^{\lambda \alpha_2} S_\omega(t) \right] \right). \end{aligned} \quad (5.11)$$

In the limit $\lambda \rightarrow 0$ we recover the linear flow

$$\Phi_{\omega,0}^t(\alpha_1, \alpha_2) = (C_\omega(t) \alpha_1 - S_\omega(t) \alpha_2, \omega S_\omega(t) \alpha_1 + C_\omega(t) \alpha_2), \quad (5.12)$$

which corresponds to the nondeformed action given by the vector field $\hat{J}_{\omega,0} = -P_2\partial_{P_1} + \omega P_1\partial_{P_2}$.

Note that although expression (5.11) has been obtained for $\omega > 0$ and $\lambda > 0$ it can be proved that it is also valid for the remaining values. In some sense, the flow has an “analytic” dependence on the parameters ω and λ which allows to extend the results obtained in a region of the parameters space to whole it.

The flow (5.11) is globally defined when $\omega \geq 0$ but is only local for $\omega < 0$ as it is easy to prove considering the integral curve passing through a point like $(0, \alpha_2)$: since the logarithm argument has to be positive one gets the inequality

$$\cosh(\lambda\alpha_2) + \sinh(\lambda\alpha_2)C_\omega(t) > 0, \quad (5.13)$$

if the product $\lambda\alpha$ is negative the definition interval of t is bounded

$$t \in (-C_\omega^{-1}(-\coth(\lambda\alpha_2)), C_\omega^{-1}(-\coth(\lambda\alpha_2))). \quad (5.14)$$

The flow (5.11) describes the action of $SO_\omega(2)$ over $T_{\lambda,2}$

$$e^{tJ} \triangleright (\alpha_1, \alpha_2) = \Phi_{\lambda,\omega}^t(\alpha_1, \alpha_2), \quad (5.15)$$

which decomposes the space $T_{\lambda,2}$ on strata of orbits depending on the values of ω :

- Case $\omega < 0$

- A stratum with only one orbit with only one point: $(0,0)$. The isotropy group is $SO_\omega(2)$.
- Four orbits determined by the points

$$\left(\frac{1-e^{-\lambda}}{\lambda\sqrt{-\omega}}, 1\right), \quad \left(\frac{1-e^\lambda}{\lambda\sqrt{-\omega}}, -1\right), \quad \left(-\frac{1-e^{-\lambda}}{\lambda\sqrt{-\omega}}, 1\right), \quad \left(-\frac{1-e^\lambda}{\lambda\sqrt{-\omega}}, -1\right). \quad (5.16)$$

These orbits are isomorphic to $\mathbb{R}$ and have the point $(0,0)$ as accumulation point.

- The remaining points of $T_{\lambda,2}$ constitute another stratum fibred by orbits diffeomorphic to $\mathbb{R}$ since they are branches of deformed hyperbolae.

- Case $\omega = 0$

- A stratum is constituted by the points $(\alpha_1, 0)$, each of them is an orbit.
- The orbits $\{(\alpha_1, \alpha_2) \mid \alpha_1 \in \mathbb{R}, \alpha_2 \neq 0 \text{ but fixed}\}$ determine a stratum.

- Case $\omega > 0$

- The point $(0,0)$ constitutes the only orbit of this stratum.
- The orbits diffeomorphic to the circle determine another stratum.

Note that the deformation associated to λ does not give qualitative changes with respect to the nondeformed case. Summarizing, we can say that the quotient spaces are isomorphic

$$T_{\lambda,2}/SO_\omega(2) \simeq T_{0,2}/SO_\omega(2). \quad (5.17)$$

5.2 Regular co-spaces

Once the flow associated to the bicrossproduct structure of $U_\lambda(\mathfrak{iso}_\omega(2))$ is known, we can study the regular co-spaces $(F_\lambda(ISO_\omega(2)), \prec, U_\lambda(\mathfrak{iso}_\omega(2)))$ and $(F_\lambda(ISO_\omega(2)), \succ, U_\lambda(\mathfrak{iso}_\omega(2)))$, which are basic elements to characterize the induced representations of $U_\lambda(\mathfrak{iso}_\omega(2))$.

The structures of both co-spaces are easily deduced combining Theorem 3.3 with expression (5.11) of the flow of $\hat{J}$. Firstly, remember that $(F_\lambda(ISO_\omega(2)))$ can be described by elements of the form

$$\phi(\alpha_1, \alpha_2), \quad \phi \in F(SO_\omega(2)), \quad (\alpha_1, \alpha_2) \in T_{\lambda,2}, \quad (5.18)$$

instead of monomials $\varphi^q a_1^r a_2^s$.

1).- For the right coregular module $(F_\lambda(ISO_\omega(2)), \prec, U_\lambda(\mathfrak{iso}_\omega(2)))$ we have that

$$[\phi(\alpha_1, \alpha_2)] \prec e^{tJ} = \phi(e^{tJ} \cdot)(\alpha_1, \alpha_2),$$

$$[\phi(\alpha_1, \alpha_2)] \prec P_1 =$$

$$\frac{\phi(2\alpha_1 \lambda e^{\lambda \alpha_2} C_\omega(\varphi) + (\omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2} - 2 \sinh(\lambda \alpha_2)) S_\omega(\varphi)}{2\lambda \cosh(\lambda \alpha_2) + \omega \lambda^3 \alpha_1^2 e^{\lambda \alpha_2} + (2\lambda \sinh(\lambda \alpha_2) - \omega \lambda^3 \alpha_1^2 e^{\lambda \alpha_2}) C_\omega(\varphi) + 2\omega \lambda^2 \alpha_1 e^{\lambda \alpha_2} S_\omega(\varphi)}(\alpha_1, \alpha_2),$$

$$[\phi(\alpha_1, \alpha_2)] \prec P_2 = \phi \frac{1}{\lambda} \ln[\cosh(\lambda \alpha_2) + \frac{1}{2} \omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2} + (\sinh(\lambda \alpha_2) - \frac{1}{2} \omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2}) C_\omega(\varphi) + \omega \lambda \alpha_1 e^{\lambda \alpha_2} S_\omega(\varphi)](\alpha_1, \alpha_2).$$

The dot stands for the argument of the function $\phi = \phi(\cdot)$. Developing in power series of t and considering the first order in the first expression and the multiplication and derivation operators associated to the basis $\varphi^q a_1^r a_2^s$ we can write the action of the generators of $U_\lambda(\mathfrak{iso}_\omega(2))$ over an arbitrary ‘function’ $f \in F_\lambda(ISO_\omega(2))$ only making the changes

$$\varphi \rightarrow \bar{\varphi}, \quad a_i \rightarrow \frac{\partial}{\partial a_i} \equiv \partial_{a_i}. \quad (5.19)$$

In this way we obtain

$$f \prec J = \partial_\varphi f,$$

$$f \prec P_1 = \frac{2\lambda \partial_{a_1} e^{\lambda \partial_{a_2}} C_\omega(\bar{\varphi}) + (\omega \lambda^2 \frac{\partial^2}{\partial a_1^2} e^{\lambda \partial_{a_2}} - 2 \sinh(\lambda \partial_{a_2})) S_\omega(\bar{\varphi})}{2\lambda \cosh(\lambda \partial_{a_2}) + \omega \lambda^3 \frac{\partial^2}{\partial a_1^2} e^{\lambda \partial_{a_2}} + (2\lambda \sinh(\lambda \partial_{a_2}) - \omega \lambda^3 \frac{\partial^2}{\partial a_1^2} e^{\lambda \partial_{a_2}}) C_\omega(\bar{\varphi}) + 2\omega \lambda^2 \partial_{a_1} e^{\lambda \partial_{a_2}} S_\omega(\bar{\varphi})} f,$$

$$f \prec P_2 = \frac{1}{\lambda} \ln[\cosh(\lambda \partial_{a_2}) + \frac{1}{2} \omega \lambda^2 \frac{\partial^2}{\partial a_1^2} e^{\lambda \partial_{a_2}} + (\sinh(\lambda \partial_{a_2}) - \frac{1}{2} \omega \lambda^2 \frac{\partial^2}{\partial a_1^2} e^{\lambda \partial_{a_2}}) C_\omega(\bar{\varphi}) + \omega \lambda \partial_{a_1} e^{\lambda \partial_{a_2}} S_\omega(\bar{\varphi})] f.$$

From the above result the action of the generator J over the ordering monomials $\varphi^q a_1^r a_2^s$ is easily evaluated

$$(\varphi^q a_1^r a_2^s) \prec J = q \varphi^{q-1} a_1^r a_2^s, \quad (5.20)$$

but it does not happen the same with P_1 and P_2 , excepting in the nondeformed case where the action reduces to

$$f \prec P_1 = \left(C_\omega(\bar{\varphi}) \frac{\partial}{\partial a_1} - S_\omega(\bar{\varphi}) \frac{\partial}{\partial a_2} \right) f, \quad f \prec P_2 = \left(\omega S_\omega(\bar{\varphi}) \frac{\partial}{\partial a_1} + C_\omega(\bar{\varphi}) \frac{\partial}{\partial a_2} \right) f. \quad (5.21)$$

So, when $\lambda = 0$ one gets:

$$\begin{aligned}(\varphi^q a_1^r a_2^s) \prec P_1 &= r\varphi^q C_\omega(\varphi) a_1^{r-1} a_2^s - s\varphi^q S_\omega(\varphi) a_1^r a_2^{s-1}, \\(\varphi^q a_1^r a_2^s) \prec P_2 &= \omega r\varphi^q S_\omega(\varphi) a_1^{r-1} a_2^s + s\varphi^q C_\omega(\varphi) a_1^r a_2^{s-1}.\end{aligned}\tag{5.22}$$

2).- The description of the left coregular module $(F_\lambda(ISO_\omega(2)), \succ, U_\lambda(\mathfrak{iso}_\omega(2)))$ is made in an analogous way. Firstly, considering Theorem 3.3 one obtains

$$\begin{aligned}e^{tJ} \succ [\phi(\alpha_1, \alpha_2)] &= \phi(\cdot e^{tJ}) \\&\left(\frac{2\lambda\alpha_1 e^{\lambda\alpha_2} C_\omega(t) + (\omega\lambda^2\alpha_1^2 e^{\lambda\alpha_2} - 2\sinh(\lambda\alpha_2)) S_\omega(t)}{2\lambda \cosh(\lambda\alpha_2) + \omega\lambda^3\alpha_1^2 e^{\lambda\alpha_2} + (2\lambda \sinh(\lambda\alpha_2) - \omega\lambda^3\alpha_1^2 e^{\lambda\alpha_2}) C_\omega(t) + 2\omega\lambda^2\alpha_1 e^{\lambda\alpha_2} S_\omega(t)} \right), \\&\frac{1}{\lambda} \ln[\cosh(\lambda\alpha_2) + \frac{1}{2}\omega\lambda^2\alpha_1^2 e^{\lambda\alpha_2} + (\sinh(\lambda\alpha_2) - \frac{1}{2}\omega\lambda^2\alpha_1^2 e^{\lambda\alpha_2}) C_\omega(t) + \omega\lambda\alpha_1 e^{\lambda\alpha_2} S_\omega(t)] \Big), \\P_1 \succ [\phi(\alpha_1, \alpha_2)] &= \alpha_1 \phi(\alpha_1, \alpha_2), \\P_2 \succ [\phi(\alpha_1, \alpha_2)] &= \alpha_2 \phi(\alpha_1, \alpha_2).\end{aligned}\tag{5.23}$$

With the same arguments that in the case of the right regular co-space the action of the generators can be written in terms of the multiplication and derivation operators

$$\begin{aligned}J \succ f &= \left[\frac{\partial}{\partial \varphi} - \bar{a}_1 \frac{1 - e^{-2\lambda \frac{\partial}{\partial a_2}}}{2\lambda} + \omega \bar{a}_2 \frac{\partial}{\partial a_1} - \frac{\lambda}{2} \omega \bar{a}_1 \frac{\partial^2}{\partial a_1^2} \right] f, \\P_i \succ f &= \frac{\partial}{\partial a_i} f, \quad i = 1, 2,\end{aligned}\tag{5.24}$$

which allows to obtain the action of the generators over the basis $\varphi^q a_1^r a_2^s$

$$\begin{aligned}J \succ (\varphi^q a_1^r a_2^s) &= q\varphi^q a_1^r a_2^{s-1} - \frac{\lambda}{2} \omega r(r-1) \varphi^q a_1^{r-1} a_2^s + \omega r \varphi^q a_1^{r-1} a_2^{s+1} \\&\quad - \frac{1}{2\lambda} \varphi^q a_1^{r+1} a_2^s + \frac{1}{2\lambda} \varphi^q a_1^{r+1} (a_2 + 2\lambda)^s \cdot s \varphi^q a_1^r a_2^{s-1}, \\P_1 \succ (\varphi^q a_1^r a_2^s) &= r \varphi^q a_1^{r-1} a_2^s, \\P_2 \succ (\varphi^q a_1^r a_2^s) &= s \varphi^q a_1^r a_2^{s-1}.\end{aligned}\tag{5.25}$$

The subalgebra $\mathcal{A}_{\omega, \lambda} = \langle a_1, a_2 \rangle$ of $F_\lambda(ISO_\omega(2))$, which is not a Hopf subalgebra, is stable under the previous action. The explicit action of the generators of $U_\lambda(\mathfrak{iso}_\omega(2))$ over a generic element $\psi(a_1, a_2)$ of $\mathcal{A}_{\omega, \lambda}$ in terms of operators adapted to the basis $a_1^r a_2^s$ is

$$\begin{aligned}J \triangleright \psi(a_1, a_2) &= \left(-\bar{a}_1 \frac{1 - e^{-2\lambda \frac{\partial}{\partial a_2}}}{2\lambda} + \omega \bar{a}_2 \frac{\partial}{\partial a_1} - \frac{\lambda}{2} \omega \bar{a}_1 \frac{\partial^2}{\partial a_1^2} \right) \psi(a_1, a_2), \\P_i \triangleright \psi(a_1, a_2) &= \frac{\partial}{\partial a_i} \psi(a_1, a_2), \quad i = 1, 2.\end{aligned}\tag{5.26}$$

The generators a_1 and a_2 of the co-space $(\mathcal{A}_{\omega,\lambda}, \triangleright, U_\lambda(\mathfrak{iso}_\omega(2)))$ verify the commutation relation

$$[a_1, a_2] = \lambda a_1 \quad (5.27)$$

defining a noncommutative geometry except in the nondeformed limit where the corresponding classical geometries are recovered.

The action of the Casimir (5.8) of $U_\lambda(\mathfrak{iso}_\omega(2))$ on $(\mathcal{A}_{\omega,\lambda}, \triangleright, U_\lambda(\mathfrak{iso}_\omega(2)))$ is given by

$$C_{\omega,\lambda} \triangleright f(a_1, a_2) = \left[\omega \frac{\partial^2}{\partial a_1^2} e^{\lambda \frac{\partial}{\partial a_2}} + \frac{4}{\lambda^2} \sinh^2\left(\frac{\lambda}{2} \frac{\partial}{\partial a_2}\right) \right] f(a_1, a_2), \quad (5.28)$$

or explicitly

$$C_{\omega,\lambda} \triangleright f(a_1, a_2) = \omega \frac{\partial^2}{\partial a_1^2} f(a_1, a_2 + \lambda) + \frac{1}{\lambda^2} [f(a_1, a_2 + \lambda) + f(a_1, a_2 - \lambda) - 2f(a_1, a_2)]. \quad (5.29)$$

This last expression shows the effect of the deformation transforming one of the derivatives on a finite difference operator.

An interesting problem is the solution of the wave equations associated to this two-parameter family of Casimir operators (in this context see, for instance Ref. [24] and references therein). Considering the group $T_{\lambda,2}$ included inside of $\mathcal{A}_{\omega,\lambda}$ by means of the exponential map, the action of the Casimir over the elements $(\alpha_1, \alpha_2) \in T_{\lambda,2}$ is given by

$$C_{\omega,\lambda} \triangleright (\alpha_1, \alpha_2) = \left[\omega \alpha_1^2 e^{\lambda \alpha_2} + \frac{4}{\lambda^2} \sinh^2\left(\frac{\lambda}{2} \alpha_2\right) \right] (\alpha_1, \alpha_2), \quad (5.30)$$

that suggests to interpret to (α_1, α_2) as a “plane wave”.

5.3 Representations of $U_\lambda(\mathfrak{iso}_\omega(2))$

The representation induced by the character of $\mathcal{L}$

$$1 \dashv (P_1^{n_1} P_2^{n_2}) = \alpha_1^{n_1} \alpha_2^{n_2} \quad (5.31)$$

(or, in other words, the representation induced by $(\alpha_1, \alpha_2) \in T_{\lambda,2}$) is obtained as follows. Theorem 3.4 gives the following result for the action of the generators in the induced representation

$$\begin{aligned} \phi \dashv J &= \phi', \\ \phi \dashv P_1 &= \phi \frac{2\lambda \alpha_1 e^{\lambda \alpha_2} C_\omega(\varphi) + (\omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2} - 2 \sinh(\lambda \alpha_2)) S_\omega(\varphi)}{2\lambda \cosh(\lambda \alpha_2) + 2\omega \lambda^3 \alpha_1^2 e^{\lambda \alpha_2} + (2\lambda \sinh(\lambda \alpha_2) - \omega \lambda^3 \alpha_1^2 e^{\lambda \alpha_2}) C_\omega(\varphi) + 2\omega \lambda^2 \alpha_1 e^{\lambda \alpha_2} S_\omega(\varphi)}, \\ \phi \dashv P_2 &= \phi \frac{1}{\lambda} \ln[\cosh(\lambda \alpha_2) + \frac{1}{2} \omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2} \\ &\quad + (\sinh(\lambda \alpha_2) - \frac{1}{2} \omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2}) C_\omega(\varphi) + \omega \lambda \alpha_1 e^{\lambda \alpha_2} S_\omega(\varphi)]. \end{aligned} \quad (5.32)$$

In the limit $\lambda \rightarrow 0$ we recover the more familiar expressions

$$\begin{aligned}\phi \dashv J &= \phi', \\ \phi \dashv P_1 &= \phi(\alpha_1 C_\omega(\varphi) - \alpha_2 S_\omega(\varphi)), \\ \phi \dashv P_2 &= \phi(\omega \alpha_1 S_\omega(\varphi) + \alpha_2 C_\omega(\varphi)).\end{aligned}\tag{5.33}$$

The local representations obtained inducing with the character of $U(\mathfrak{so}_\omega(2))$ given by

$$J^m \vdash 1 = c^m,\tag{5.34}$$

are

$$\begin{aligned}e^{tJ} \vdash (\alpha_1, \alpha_2) &= e^{tc} \\ &\times \left(\frac{2\lambda \alpha_1 e^{\lambda \alpha_2} C_\omega(t) + (\omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2} - 2 \sinh(\lambda \alpha_2)) S_\omega(t)}{2\lambda \cosh(\lambda \alpha_2) + \omega \lambda^3 \alpha_1^2 e^{\lambda \alpha_2} + (2\lambda \sinh(\lambda \alpha_2) - \omega \lambda^3 \alpha_1^2 e^{\lambda \alpha_2}) C_\omega(t) + 2\omega \lambda^2 \alpha_1 e^{\lambda \alpha_2} S_\omega(t)}, \right. \\ &\left. \frac{1}{\lambda} \ln[\cosh(\lambda \alpha_2) + \frac{1}{2} \omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2} + (\sinh(\lambda \alpha_2) - \frac{1}{2} \omega \lambda^2 \alpha_1^2 e^{\lambda \alpha_2}) C_\omega(t) + \omega \lambda \alpha_1 e^{\lambda \alpha_2} S_\omega(t)] \right),\end{aligned}$$

$$P_i \vdash (\alpha_1, \alpha_2) = \alpha_i(\alpha_1, \alpha_2), \quad i = 1, 2,\tag{5.35}$$

which can be written in an equivalent way using an arbitrary element, $\psi(a_1, a_2)$, of $U(\mathfrak{t}_{\lambda,2})$, as

$$\begin{aligned}J \vdash \psi(a_1, a_2) &= \left[c - \bar{a}_1 \frac{1 - e^{-2\lambda \frac{\partial}{\partial a_2}}}{2\lambda} + \omega \bar{a}_2 \frac{\partial}{\partial a_1} - \frac{\lambda}{2} \omega \bar{a}_1 \frac{\partial^2}{\partial a_1^2} \right] \psi(a_1, a_2), \\ P_i \vdash \psi(a_1, a_2) &= \frac{\partial}{\partial a_i} \psi(a_1, a_2), \quad i = 1, 2.\end{aligned}\tag{5.36}$$

Notice that if we take $c = 0$, which is equivalent to consider the character determined by the count of $U(\mathfrak{so}_\omega(2))$, the action (5.26) of the co-space $(\mathcal{A}_{\omega,\lambda}, \triangleright, U_\lambda(\mathfrak{iso}_\omega(2)))$ is recovered.

The Casimir action for the local representation coincides with those given by (5.28) since only the generators P_i are presented.

It is worthy to note that also in this case in the limit $\lambda \rightarrow 0$ we obtain

$$\begin{aligned}J \vdash \psi(a_1, a_2) &= \left[c - a_1 \frac{\partial}{\partial a_2} + \omega a_2 \frac{\partial}{\partial a_1} \right] \psi(a_1, a_2), \\ P_i \vdash \psi(a_1, a_2) &= \frac{\partial}{\partial a_i} \psi(a_1, a_2), \quad i = 1, 2\end{aligned}\tag{5.37}$$

that it is in agreement with the results for local representations [25].

6 Concluding remarks

The induction procedure that we have used here is not, strictly speaking, a generalization of Mackey's induction method for Lie groups. The concept of co-space generalizes in an algebraical

way the concept of G -space (being G a transformation group), and is connected with the induced representations in some sense similar to the nondeformed case. In Ref. [6] we presented a more general method that, obviously, can also be used for bicrossproduct algebras. However, it is necessary the knowledge of pairs of dual basis of the corresponding Hopf algebra and its dual, but here this requirement is not necessary in the method used in this work.

We have made use of the vector fields to compute commutators. For bicrossproduct Hopf algebras, like used in this work $H = K \bowtie \mathcal{L}$, there is a connection between the representations of H induced by characters of L and one-parameter flows. This relation allows to associate to quantum bicrossproduct groups dynamical systems. Here we have only sketched this situation that will be analyzed with more detail in a forthcoming paper.

The equations associated to the Casimir operators, as in the nondeformed case, will give the behaviour of the ‘deformed’ quantum systems. A procedure for this solution can be found in [24]. Note that q -special polynomials and q -functions may appear as solutions of these q -Casimir equations.

As we mention in Section 2 the bicrossproduct structure gives a $*$ -structure for which the representations are, essentially, unitary. The equivalence of the induced representations is given by Theorem 3.4, which establishes a correspondence among classes of induced representations and orbits of L under the action of K . This result is analogous to the Kirillov orbits method [26]. The problem of the irreducibility of the representations is still open. Partial results for particular cases have been obtained (see Ref. [4] for the $(1+1)$ κ -Galilei algebra and [22] for the quantum extended $(1+1)$ Galilei algebra).

Acknowledgments

This work has been partially supported by DGES of the Ministerio de Educación y Cultura de España under Project PB98-0360, and the Junta de Castilla y León (Spain).

References

- [1] O. Arratia and M.A. del Olmo, *Representations of Quantum Bicrossproduct Algebras*, preprint UVA 2001; math.QA/0110275.
- [2] E. P. Wigner, *Ann. Math.* **40**, (1939), 149.
- [3] J.F. Cariñena, M.A. del Olmo and M. Santander, *J. Phys. A* **17**, (1984) 3091.
- [4] O. Arratia and M.A. del Olmo, *Induced representations of quantum groups* in *Anales de Física, Monografías* vol. 5 (Ciemat/RSEF, Madrid 1998); math.QA/0110265.
- [5] O. Arratia and M.A. del Olmo, *Elements of the theory of induced representations for quantum groups*, *Publicaciones de la RSME* vol. 1, (RSME, Madrid 2000); math.QA/0110266.
- [6] O. Arratia and M.A. del Olmo, *J. Math. Phys.* **41**, (2000) 4817.

- [7] S. Majid, *Class. Quant. Grav.* **5**, (1988) 1587; *J. Alg.* **130**, (1990) 17; *Pac. J. Math.* **141**, (1990) 311; *Isr. J. Math.* **72**, (1990) 133.
- [8] S. Majid, *Foundations of quantum group theory* (Cambridge Univ. Press, Cambridge 1995).
- [9] A. Connes, *Non-commutative geometry*. Academic Press, London 1994.
- [10] G.V. Mackey, *Unitary groups representations* (Benjamin/Cummings, Reading, 1978).
- [11] M. Santander, F.J. Herranz and M.A. del Olmo, *Kinematics and homogeneous spaces for symmetrical contractions of orthogonal groups* in “Group Theoretical Methods in Physics”, pp. 455. Eds. M.A. del Olmo, M. Santander and J. Mateos-Guilarte, Anales de Física, Monografías, 1 vol. I. (CIEMAT/RSEF, Madrid 1993).
- [12] A. Ballesteros, F. J. Herranz, M. A. del Olmo and M. Santander, *J. Phys. A: Math. Gen.* **26** (1993), 5801.
- [13] A. Ballesteros, F. J. Herranz, M. A. del Olmo and M. Santander, *J. Phys. A: Math. Gen.* **27** (1994), 1283.
- [14] A. Ballesteros, F. J. Herranz, M. A. del Olmo and M. Santander, *J. Math. Phys.* **35** (1994), 4928.
- [15] A. Ballesteros, F. J. Herranz, M. A. del Olmo and M. Santander, *Lett. Math. Phys.* **33** (1995), 273.
- [16] J.A. de Azcárraga, M.A. del Olmo, J.C. Pérez-Bueno and M. Santander, *J. Phys. A: Math. Gen.* **30** (1997), 3069.
- [17] V.K. Dobrev, *J. Phys. A* **27**, (1994) 4841 and 6633; *Induced representations and invariant operators for quantum groups*, to appear in *Turk. J. Phys.*
- [18] A. González-Ruiz and L.A. Ibort, *Phys. Lett.* **B296**, (1992) 104.
- [19] N. Ciccoli, *Induction of quantum group representations*, math/9804138.
- [20] P. Maślanka, *J. Math. Phys.* **35**, (1994) 5047.
- [21] S. Giller, C. Gonera, P. Kosiński and P. Maślanka, hep-th 9505007.
- [22] F. Bonechi, R. Giachetti, E. Sorace and M. Tarlini, *Lett. Math. Phys.* **43**, (1998) 309.
- [23] V. Chari and A. Pressley, *A guide to quantum groups* (Cambridge Univ. Press, Cambridge 1994).
- [24] F. Bonechi, E. Celeghini, R. Giachetti, M.A. del Olmo, E. Sorace and M. Tarlini, *J. Phys. A: Math. Gen.* **29**, (1996) 7973.
- [25] M. Gadella, M.A. Martín, L.M. Nieto and M.A. del Olmo, *J. Math. Phys.* **32** (1992) 1182.
- [26] A.A. Kirillov, *Elements of the Theory of Representations* (Springer, Berlin 1976).